

**Officers :**

G. H. HICKS, President, Owosso, Mich. *Jt Collier*  
E. L. BYINGTON, A. M., Gen. Sec'y, Boulder, Col.  
MISS H. H. KEYES, Treasurer, Watertown, N. Y.  
A. B. SIMONDS, Ex. Council, 30 Thayer Hall, Cambridge, Mass.  
J. M. HOLZINGER, A. B., Executive Council, Winona, Minn.

**Division Secretaries :**

A—MISS C. G. DU BOIS, 9 Hillside Ave., Waterbury, Conn.  
B—C. H. EHRENFIELD, A. M., York, Pa.  
C—CHAS. A. DAVIS, S. B., Alma, Mich.  
D—J. M. HOLZINGER, A. B., Winona, Minn.  
E—E. R. DREW, 472 24th St, Oakland, Calif.

GRAY MEMORIAL BOTANICAL CHAPTER,

—OF THE—

AGASSIZ ASSOCIATION.

Owosso Mich July 2/90

Dr Sereno Watson, Dear Sir:-

I take the liberty of sending you by this mail a ph of 110 or 110 plants collected by myself & wife in June 1890, at Veta Pass, Colorado, 206 miles southwest of Denver, in the Sangre de Christo range, at an altitude of 9370 ft. or more.

Regret that the season was not late enough to admit of collecting much fruit, rendering identification more difficult to me, not being at all acquainted with the flora of Colorado.

Perhaps you will find some of the plants interesting, possibly one or two new varieties.

In No 6 - (*Clematis Douglasii* Hook ?) the leaflets are quite variable and the sepals deep purple without & greenish white within contrary to description in Coulter's Manual

The plant often grows in clumps varying from 8 to 14 inches high - Very common on open mountain sides, presenting striking appearance.

No 8 - *Castilleja* Drummondii? puzzled me from the <sup>invariate</sup> drooping position of the pods

No 19 Battystis palustris var angustifolius?  
was the most common begoniacous plant in the vicinity, being abundant in light woods. Short, generally not over a foot high - flowers rather small & light colored, often nearly white, with light purple or rose-colored lines - Stem twisted - rachis sometimes tendril bearing especially in taller specimens - no fruit obtained -

No 52. *Geranium* — ? was quite common in light woods & on shady banks of mountain brooks. Flowers light purple often nearly white -

"differs from *G. Carolinianum* in elongated peduncle, not branched from base - lobes of lowest radical leaves blunt - petals generally almost white, with pale rose-colored lines - scarcely or not at all emarginate -

**Officers:**

G. H. HICKS, President, Owosso, Mich.  
E. L. BYINGTON, A. M., Gen. Sec'y, Boulder, Col.  
MISS H. H. KEYES, Treasurer, Watertown, N. Y.  
A. B. SIMONDS, Ex. Council, 30 Thayer Hall, Cambridge, Mass.  
J. M. HOLZINGER, A. B., Executive Council, Winona, Minn.

**Division Secretaries:**

A—MISS C. G. DU BOIS, 9 Hillside Ave., Waterbury, Conn.  
B—C. H. EHRENFIELD, A. M., York, Pa.  
C—CHAS. A. DAVIS, S. B., Alma, Mich.  
D—J. M. HOLZINGER, A. B., Winona, Minn.  
E—E. R. DREW, 472 24th St. Oakland, Calif.

GRAY MEMORIAL BOTANICAL CHAPTER,

—OF THE—

AGASSIZ ASSOCIATION.

If you wish I will send printed labels  
after receiving list of names & you can keep  
them for the College Herb. if thought  
worthy.

If you are too busy to look them  
over now you will kindly notify me &  
I will send stamps to bear the  
forwarded elsewhere.

Hoping to hear from you soon & to  
be excused for intruding upon your  
time, I am  
Yours truly

G. H. Hicks

If you wish for extra spec. of any  
number please notify me.

Hicks